

Sterols from Margarine by FID on ZB-MR2

Column: Zebtron™ ZB-MultiResidue™-2, GC Cap. Column 30 m x 0.25 mm x 0.20 μm, Ea

Phase: Proprietary Pesticides Phase

Dimensions: 30 meters x 0.25 mm x 0.2 μm

Order No: 7HG-G017-10

Oven Profile: 220 °C to 320 °C @ 13 °C/min.

Carrier Gas: Constant Flow Helium, 2 mL/min


Injection: Splitless :1 0.5 μL @ 300°C

Detection: Flame Ionization (FID) (300°C)

Analyst Note: Sample Prep: Saponification, Solid Phase Extraction (SPE), and Derivatization.



Products used in this application:



ANALYTES:

- 1 Campesterol
- 2 Stigmasterol
- 3 b-Sitosterol
- 4 Betulin (internal standard)



Sample Preparation Details

for GC Application ID No.: 16756

Sterols from Margarine by FID on ZB-MR2

PRODUCT DESCRIPTION:

Strata® C18-E (55 µm, 70 Å), 1 g / 6 mL, Tubes , 30/Pk

Order No.: 8B-S001-JCH

SOLID PHASE EXTRACTION (SPE) PROCEDURE:

Note: The solvent volumes shown below are for a 1 g bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:

Load:

Saponify sample in 10 mL 0.5 M KOH in ethanol at 77 °C for 20 min.

When cool, add 10 mL of 0.22 mg/ml betulin (istd) in chloroform and shake.

Adjust 3 mL aliquot of sample to pH 2.5 with 5 M HCl.

Syringe filter aliquot through Phenex 30 mm 0.45 µm teflon filter.

Purify via Strata C18-E (55 µm, 70 Å).

[Condition SPE with 5 mL methanol then 5 mL chloroform.

Load 1 mL of filtrate sample.

Elute with 15 mL 5% methanol in chloroform.]

Blow down eluate to 1 mL.

Transfer 200 µL to 500 µL insert and dry under nitrogen.

Reconstitute with 100µL pyridine and 100 µL BSTFA:TMCS 99:1 and derivatize at 70°C for 20 min.

Wash:

Dry:

Elute:

Final Prep and Analysis:

Initial Sample Weight / Unit: 0.70 grams

Inject: 0.5 µL on HPLC Flame Ionization (FID) @ (300°C)

ANALYTES:	Spiked Conc. (ng/mL)	Log P	pKa	% Rec	%RSC (n=0)
1 Campesterol	0				
2 Stigmasterol	0				
3 b-Sitosterol	0				
4 Betulin	0				

Note: This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2019 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at info@phenomenex.com



Phenomenex products are available worldwide.

www.phenomenex.com

info@phenomenex.com