

PEGylated L-Chymotrypsinogen A (N-Terminal PEG 20KDa) on BioSep2000 (2)

Column: BioSep™ 5 µm SEC-s2000 145 Å, LC Column 300 x 7.8 mm, Ea

Dimensions: 300 x 7.8 mm ID

Order No: 00H-2145-K0

Elution Type: Isocratic

Eluent A: 100mM Phosphate buffer pH 6.8□□

Gradient Profile:	Step No.	Time (min)	Pct A
	1	0	100


Flow Rate: 1 mL/min

Col. Temp.: ambient


Detection: UV-Vis Abs.-Variable Wave.(UV) @ 220 nm (ambient)

Analyst Note: Application Topic: Monitoring protein PEGylation and purifying PEGylated proteins from their reaction

Therapeutic proteins are often PEGylated to increase their serum lifetime; however, such reactions typically generate a heterogeneous product that can be difficult to characterize and purify. PEGylated proteins are usually purified by GFC or RPC after synthesis and the PEGylation reaction is often monitored using SDS-PAGE. In this example application, protein was reacted using N-terminal favoring conditions (phosphate pH 6.5 with cyanoborohydrate and 5% PEG-aldehyde excess). For reaction monitoring, intact protein and reaction timepoints at 0 minutes, 30 minutes, 60 minutes and an overnight reaction



Products used in this application:



ANALYTES:

- 1 4 PEG / Chymo A complex (co-elution)
- 2 3 PEG / Chymo A complex (co-elution)
- 3 2 PEG / Chymo A complex (co-elution)
- 4 PEGylated Chymotrypsinogen A
- 5 Chymotrypsinogen A

